

The Peculiarity of a Negative Coordinate Axis in Dyonic Solutions of Noncommutative $\mathcal{N} = 4$ Super Yang-Mills

Aalok Misra

Harish Chandra Research Institute,
Jhusi, Allahabad - 211 019, India
email: aalok@mri.ernet.in

Abstract

We show that in the neighborhood of a negative coordinate axis, the $U(1)$ sector of the static dyonic solutions to the noncommutative $U(4)$ $\mathcal{N} = 4$ Super Yang-Mills (SYM) can be consistently decoupled from the $SU(4)$ to *all orders in the noncommutativity parameter*. We show the above decoupling in two ways. First, we show the noncommutative dyon being the same as the commutative dyon, is a consistent solution to noncommutative equations of motion in the abovementioned region of noncommutative space. Second, as an example of decoupling of a non-null $U(1)$ sector, we also obtain a family of solutions with nontrivial $U(1)$ components for all components of the gauge field in the same region of noncommutative space. In both cases, the $SU(4)$ and $U(1)$ components separately satisfy the equations of motion.

1 Introduction

Nonperturbative solutions in Super Yang-Mills (SYM) theories have been an important area of work. More recently, static dyonic solutions in $SU(N)$, $N \geq 3, 4$ $\mathcal{N} = 4$ SYM corresponding to planar and non-planar (respectively) string junctions have been obtained in [1, 2] (See also references therein). Also, solitonic solutions in noncommutative SYM have been considered by others before (See [3, 4] and references therein). Now, the gauge group $SU(N)$, in general is not allowed in noncommutative gauge theories [5] (See [6] though, where the gauge parameter in noncommutative gauge transformations is allowed to depend on the usual $SU(N)$ gauge parameter and gauge fields. The explicit form of the map to all orders in the noncommutativity parameter [which is what we are interested in, in this paper] is quite involved and has not been worked out.). One of the questions that we address in this work is whether it is possible to construct solutions to the equations of motion of a noncommutative supersymmetric gauge theory in which one can decouple the $U(1)$ components from

the SU components of the fields at least in some region of the noncommutative space, *to all orders in the noncommutativity parameter*. There are the following two additional motivations for this work. The equality of the angular momentum and (null) quadrupole moment of the commutative and noncommutative monopoles of (S)U(2) $N = 2$ SYM up to $O(\theta^2)$ was shown in [13] and in all of space. This makes it natural to pose the following question: is it possible to construct solutions of a noncommutative supersymmetric gauge theory that are characterized by quantities that receive no noncommutativity corrections *to all orders in the noncommutativity parameter*? Finally, it is of obvious interest to construct solutions that are valid up to all orders in the noncommutativity parameter.

In this paper, we show that for noncommutative $U(4)$ $\mathcal{N} = 4$ SYM, the static dyonic solutions in the neighborhood of a negative coordinate axis of the noncommutative space, are such that one can decouple the $U(1)$ from the $SU(4)$, and construct a 6-parameter family of noncommutative dyonic solutions (for a non-null $U(1)$ decoupling). Even though, the decoupling has been shown for a very limited region of the noncommutative space, the reason the solutions are of interest is because they have the same electric and magnetic charges as and energy differing infinitesimally from the commutative dyonic solutions of [1], *to all orders in the noncommutativity parameter*. This points to the peculiar nature of the the negative x^3 axis when constructing dyonic solutions to noncommutative $\mathcal{N} = 4$ SYM, which requires further investigation.

The paper is organized as follows. In section 2, we discuss the basics needed for getting static dyonic solutions in commutative $\mathcal{N} = 4$ Super Yang-Mills (SYM). In section 3, we discuss the modifications in the techniques of section 2 to get static dyonic solutions of noncommutative $\mathcal{N}=4$ SYM. It has two subsections: (a short) 3.1 on decoupling of a null $U(1)$ from the $SU(4)$, and 3.2 on decoupling of a non-null $U(1)$ from the $SU(4)$. In section 4, we give a summary of and a discussion on the results obtained in the paper, and indicate possible future directions for work.

2 Basics

In this section, we give a brief review of the techniques given in [1] to obtain dyonic solutions in commutative $SU(4)$ $\mathcal{N} = 4$ SYM.

It is advisable to write the equations of motion as if one were working in curved space-time. The reason is that for solutions of the equations of motion, it is convenient to go to the stereographic coordinate system (See Fig 1: $z = e^{i\phi} \tan \frac{\theta}{2}$) in which the metric is not constant. The equations of motion for commutative $SU(4)$

$\mathcal{N} = 4$ SYM are given by:

$$\begin{aligned} D_i(\sqrt{g}g^{ii_1}g^{jj_1}F_{i_1j_1}) &= i\sqrt{g}\sum_J[D^j\Phi^J, \Phi^J]; \\ D_i(\sqrt{g}g^{ij}D_j\Phi^J) &= \sqrt{g}[\Phi^I, [\Phi^I, \Phi^J]]. \end{aligned} \quad (1)$$

The component-form of the above equations is given in [1] which we won't be repeating here. Following the close parallel with Skyrme models[7], one uses the harmonic-map ansatz to solve the equations of motion - the Higgs and gauge fields are assumed to have the following form:

$$\begin{aligned} A_z &= A_z^a T^a = -(1 + c_1 - c_2 - c_0)(r)(\partial P_0)P_0 + (c_2 - c_1)(r)[P_1, \partial P_1] + (1 - c_2(r))[P_2, \partial P_2], \\ A_r &= 0; \\ \Phi^I &= \Phi^{I,a} T^a = (b_2^I + b_0^I + b_1^I)(r)(P_0 - \frac{1}{4}\mathbf{1}_4) + (b_2^I + b_1^I)(r)(P_1 - \frac{1}{4}\mathbf{1}_4) + b_2^I(r)(P_2 - \frac{1}{4}\mathbf{1}_4), \end{aligned} \quad (2)$$

T^a being the generators of $SU(4)$, I indexing the number of scalars that are turned on, and for this paper taking values 1,2 and 3, $c_{0,1,2}(r)$ and $b_{0,1,2}^I(r)$ being the 'profile functions' of the gauge field and Higgs respectively, and P_i the orthogonal set of $\mathbf{CP}^3$ sigma model projectors defined as:

$$\begin{aligned} P_0 &= \frac{ff^\dagger}{f^\dagger f}; \\ P_j &= \frac{(\Delta^j f)(\Delta^j f)^\dagger}{|\Delta^j f|^2}, \quad j = 1, 2, \end{aligned} \quad (3)$$

where

$$\begin{aligned} \Delta f &\equiv \partial f - f \frac{f^\dagger \partial f}{|f|^2}; \\ f &\equiv (1, \sqrt{3}z, \sqrt{3}z^2, z^3)^t, \end{aligned} \quad (4)$$

and $\Delta^2 f = \Delta(\Delta f)$. They can be thought of as hermitian maps from S^2 to $\mathbf{CP}^3$. The explicit forms of the projectors P_i s can be worked out using Mathematica or Maple.

The substitution of the harmonic map ansatz into the equations of motion (written in components) either results in an equation of motion being identically satisfied, or a system of non-linear second-order coupled differential equations involving the Higgs and gauge field profile functions that can be solved using techniques referred to in [1]. The solutions corresponding to (non-)planar string networks (for non-planar, assuming an infinitesimal non-planarity corresponding to an infinitesimal deformation from the BPS nature of the planar solution) were then obtained in [1].

3 Noncommutative dyonic solutions

In this section we derive a class of static noncommutative dyonic solutions assuming $r - z$ and $r - \bar{z}$ noncommutativity only. Thus, the star product of two functions $f(r, z, \bar{z})$ and $h(r, z, \bar{z})$ will be:

$$f(r, z, \bar{z}) * h(r, z, \bar{z}) = e^{\frac{i}{2}[\Theta^{rz}(\partial_r^1 \partial^2 - \partial_r^2 \partial^1) + \Theta^{r\bar{z}}(\partial_r^1 \bar{\partial}^2 - \partial_r^2 \bar{\partial}^1)]} f(r_1, z_1, \bar{z}_1) h(r_2, z_2, \bar{z}_2)|_{1=2}. \quad (5)$$

We show that using stereographic coordinates, in the neighborhood of the South Pole, or equivalently the neighborhood of the negative x^3 axis, one can consistently decouple in the equations of motion, the $U(1)$ from the $SU(4)$ components of the Higgs and the gauge fields.

The noncommutative equations of motion are given by:

$$\begin{aligned} \hat{D}_i * (\sqrt{g} * g^{ii_1} * g^{jj_1} * \hat{F}_{i_1 j_1}) &= i\sqrt{g} * \sum_J [\hat{D}^j * \hat{\Phi}^J, \hat{\Phi}^J]_*; \\ \hat{D}_i * (\sqrt{g} * g^{ij} * \hat{D}_j * \hat{\Phi}^J) &= \sqrt{g} * [\hat{\Phi}^I, [\hat{\Phi}^I, \hat{\Phi}^J]_*]_*. \end{aligned} \quad (6)$$

As we are considering only $r - z, r - \bar{z}$ -noncommutativity, the projectors of [1] are still valid here. We assume the following ansatz for the gauge field $\hat{A}_z, \hat{A}_r$ and the Higgs field $\hat{\Phi}^I$:

$$\begin{aligned} \hat{A}_z &= \hat{A}_z^a T^a + i\epsilon F_z(\Theta) g_z(r) \hat{\mathcal{A}}_z^0 \mathbf{1}_4; \\ \hat{A}_r &= i\epsilon F_r(\Theta) g_r(r) \hat{\mathcal{A}}_r^0 \mathbf{1}_4; \\ \hat{\Phi}^I &= \hat{\Phi}^{I,a} T^a + i\epsilon F_\Phi(\Theta) g_\Phi^I(r) \hat{\phi}^0 \mathbf{1}_4, \end{aligned} \quad (7)$$

where $F_{z,r,\Phi}(\Theta)$ are functions of the noncommutativity parameter Θ which vanish as Θ is set to zero and are bounded as $\Theta \rightarrow \infty$, and

$$\begin{aligned} \hat{A}_z^a T^a &= -(1 + c_1 - c_2 - c_0)(r) * (\partial P_0) P_0 + (c_2 - c_1)(r) * [P_1, \partial P_1] + (1 - c_2(r)) * [P_2, \partial P_2], \\ \hat{\Phi}^{I,a} T^a &= (b_2^I + b_0^I + b_1^I)(r) * (P_0 - \frac{1}{4} \mathbf{1}_4) + (b_2^I + b_1^I)(r) * (P_1 - \frac{1}{4} \mathbf{1}_4) + b_2^I(r) * (P_2 - \frac{1}{4} \mathbf{1}_4), \end{aligned} \quad (8)$$

$c_{0,1,2}(r)$ and $b_{0,1,2}^I(r)$ being the ‘profile functions’ of the gauge field and Higgs respectively as in (2). We have thus taken an ansatz in which the $U(4)$ fields have infinitesimal $U(1)$ components. We will work up to $O(\epsilon)$. Hence, the noncommutative $U(1)$ field strengths are the same as the commutative $U(1)$ field strengths.

There is no star product in the ansatz (7) in the $U(1)$ components of the fields because as argued below, the star products can be dropped in the neighborhood of the South Pole in the $SU(4)$ -components of the fields, and we assume similar behavior for the $U(1)$ and $SU(4)$ components. The ansatz for the $U(1)$ components that we choose in (10) below, is such that if one were to put in star product between the radial and angular functions, then it (the star product) can not be dropped in the neighborhood of the South Pole.

As *expanding the star product in (8) will involve taking derivatives of $(\partial)P_i$* , using the explicit forms of the projectors that can be worked out using Mathematica or Maple, one sees that *one can drop $(\partial)\partial P_i$ relative to $(\partial)P_i$ as one approaches the neighborhood of the South Pole*, as the former is suppressed relative to the latter by terms of $O(\frac{1}{|z|^m})$, $m > 0$ that vanish as $|z| \rightarrow \infty$ (i.e. in the neighborhood of the south pole). *One thus has for the SU part of $\hat{A}_z, \hat{\Phi}$ the same expression as in [1].* One can argue similarly that the $*$ products in (6) can also be dropped (in the neighborhood of the South Pole). We now discuss the decoupling from $SU(4)$, of a null $U(1)$ in **3.1** and non-null $U(1)$ in **3.2**.

3.1 Decoupling of null $U(1)$

Using (7), for null $U(1)$ components, one can require consistently that the $SU(4)$ and $U(1)$ components, separately, satisfy the equations of motion, and one can show there is no mixing between the $SU(4)$ and $U(1)$ sectors. Hence, one can consistently set the $U(1)$ components to zero. The entire analysis of [1] to show the connection between (non-)planar stable (non-)BPS string junctions and dyon solutions, then goes through even in the neighborhood of the South Pole of the abovementioned noncommutative space if one notices and uses the following transformation that takes one from the North Pole to the South Pole (and is also a symmetry of the equations of motion of the $SU(4)$ part of the solutions, as given in reference two of [1]):

$$c_0 \leftrightarrow c_2; \quad b_0^I \leftrightarrow b_2^I. \quad (9)$$

One also uses the fact that if $\hat{\Phi}$ is a solution then $-\hat{\Phi}$ is also a solution to the equations of motion.

3.2 Decoupling of non-null $U(1)$

Here we discuss in detail, the decoupling of non-null $U(1)$ components of the Higgs and the gauge fields from their $SU(4)$ components in the neighborhood of the South Pole of the noncommutative space. To show the decoupling of a non-null $U(1)$ from

$SU(4)$, we choose the following ansatz for the $U(1)$ components (of the Higgs and gauge fields):

$$\begin{aligned}\hat{\phi}^0, \hat{\mathcal{A}}_z^0 &= \text{Det}[(P_0 - a_{\Phi,z}\mathbf{1}_4)(P_1 - M_{\Phi,z})(P_2 - M_{\Phi,z})] + \text{cc} \\ \hat{\mathcal{A}}_r^0 &= z\text{Det}[(P_0 - a_r\mathbf{1}_4)(P_1 - M_r)(P_2 - M_r)] + \text{cc}.\end{aligned}\tag{10}$$

where

$$M_i = \begin{pmatrix} 0 & b_i & 0 & 0 \\ c_i & 0 & 0 & 0 \\ 0 & 0 & 0 & e_i \\ 0 & 0 & f_i & 0 \end{pmatrix},\tag{11}$$

where $i \equiv r, z, \Phi$, and the parameters a_i, b_i, c_i, e_i and f_i have been introduced to get non-zero determinants using the projectors and some constant matrices; the parameters get constrained later. We now assume that one approaches the South pole along any path very close to $\phi(\equiv \text{azimuthal angle}) = \frac{\pi}{2}$. This implies that $\text{Re}z$ is finite, but $\text{Im}z$ approaches infinity. Hence, one sees that $\partial\bar{\partial}\hat{\mathcal{A}}_r^0$ (as shown in (A5)) and $(\partial - \bar{\partial})(\hat{\phi}^0, \hat{\mathcal{A}}_z^0)$ (as shown in (A8)) are real. If

$$b_i = f_i, \quad c_i = e_i,\tag{12}$$

then $\partial\left((1 + |z|^2)^2(\partial + \bar{\partial})(\hat{\phi}^0, \hat{\mathcal{A}}_z^0)\right)$ (as shown in (A9)) will be finite and real as $\text{Im} z \rightarrow \infty$. These considerations become useful when one solves the equations of motion for the $U(1)$ components as done in (19) and (20) in the neighborhood of the South Pole.

From power (of $|z|$ or $\text{Im}z$) counting arguments using expressions given in the appendix, one can require consistently that the $SU(4)$ and $U(1)$ components, separately, satisfy the equations of motion, and one can show there is no mixing between the $SU(4)$ and $U(1)$ sectors. As the decoupling for non-null $U(1)$ is non-trivial, to illustrate the point, we consider the example of $\hat{F}_{rz}$. We see that up to $O(\epsilon)$:

$$\hat{F}_{rz}^{U(4)} = \hat{F}_{rz}^{SU(4)} + \hat{F}_{rz}^{U(1)} - i[\hat{A}_r^{U(1)}, \hat{A}_z^{SU(4)}]_*,\tag{13}$$

Now,

$$\begin{aligned}[\hat{A}_r^{U(1)}, \hat{A}_z^{SU(4)}]_* &= i\Theta^{rz}\left(\partial_r\hat{A}_r^{U(1)}\partial\hat{A}_z^{SU(4)} - \partial\hat{A}_r^{U(1)}\partial_r\hat{A}_z^{SU(4)}\right) \\ &+ i\Theta^{r\bar{z}}\left(\partial_r\hat{A}_r^{U(1)}\bar{\partial}\hat{A}_z^{SU(4)} - \bar{\partial}\hat{A}_r^{U(1)}\partial_r\hat{A}_z^{SU(4)}\right) + O(\Theta^2).\end{aligned}\tag{14}$$

Using expressions in the appendix, one sees that:

$$\begin{aligned}
\partial_r \hat{A}_r^{U(1)} &\equiv O(1), \quad \partial \hat{A}_r^{U(1)}, \bar{\partial} \hat{A}_r^{U(1)} \equiv O(1); \\
\partial \hat{A}_z^{SU(4)} &\equiv O\left(\frac{1}{|z|^3}\right) \times M, \quad \bar{\partial} \hat{A}_z^{SU(4)} \equiv O\left(\frac{1}{|z|^4}\right) \times \mathbf{1}_4, \quad \partial_r \hat{A}_z^{SU(4)} \equiv O\left(\frac{1}{|z|^2}\right) \times M; \\
\partial_r \hat{A}_z^{U(1)} &\equiv O(1), \quad \partial \hat{A}_z^{U(1)}, \bar{\partial} \hat{A}_z^{U(1)} \equiv O\left(\frac{1}{|z|^2}\right) (\text{relevant for } [\hat{A}_r^{U(1)}, \hat{A}_z^{U(1)}]_*),
\end{aligned} \tag{15}$$

where

$$M \equiv \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \tag{16}$$

Thus

$$\hat{F}_{rz}^{U(1)} \equiv O(1); \hat{F}_{rz}^{SU(4)} \equiv O\left(\frac{1}{|z|^2}\right) \times M, \tag{17}$$

and in Table 1, we show which terms can be dropped relative to which terms as $Imz \rightarrow \infty$. Similar reasoning can be used to argue that terms of higher order in Θ ,

Table 1: Consequences of power counting done in (15)

Term that can be dropped	relative to
$\partial_r \hat{A}_r^{U(1)} \partial \hat{A}_z^{SU(4)}$	$\hat{F}_{rz}^{SU(4)}$
$\partial \hat{A}_r^{U(1)} \partial_r \hat{A}_z^{SU(4)}$	$\hat{F}_{rz}^{SU(4)}$
$\partial_r \hat{A}_r^{U(1)} \bar{\partial} \hat{A}_z^{SU(4)}$	$\hat{F}_{rz}^{U(1)}$
$\bar{\partial} \hat{A}_r^{U(1)} \partial_r \hat{A}_z^{SU(4)}$	$\hat{F}_{rz}^{SU(4)}$

can also be dropped relative to the commutative counterparts. Hence, as $Imz \rightarrow \infty$,

$$\hat{F}_{rz}^{U(4)} \sim \hat{F}_{rz}^{SU(4)} + \hat{F}_{rz}^{U(1)}, \tag{18}$$

showing the stated decoupling.

By requiring that the SU components of the Higgs and gauge fields still satisfy the commutative equations of motion (having dropped the star product at the South pole), the $U(1)$ components should satisfy these equations:

$$\frac{dg_z}{dr} (\partial - \bar{\partial}) \hat{\mathcal{A}}_z^0 - 2g_r \partial \bar{\partial} \hat{\mathcal{A}}_r^0 = 0;$$

$$\begin{aligned} \frac{d^2 g_z}{dr^2} \hat{\mathcal{A}}_z^0 - g_r \partial \hat{\mathcal{A}}_r^0 + \frac{g_z}{2r^2} \partial \left((1 + |z|^2)^2 (\partial + \bar{\partial}) \hat{A}_z \right) &= 0 \\ \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dg_\Phi^I}{dr} \right) \hat{\phi}^0 + \frac{(1 + |z|^2)^2}{r^2} g_\Phi^I \partial \bar{\partial} \hat{\phi}^0 &= 0. \end{aligned} \quad (19)$$

The $*$ product has been dropped as it will generate $O(\frac{1}{|z|^m})$ - corrections (relative to $O(1)$), which can be dropped in the neighborhood of the south pole. We now discuss a family of dyonic solutions to (19).

One gets (setting $c_r = e_r$ and $b_r = f_r$ for convenience):

$$\begin{aligned} \frac{dg_z}{dr} + 2g_r \frac{[(b_r c_r)^3 (2c_r - b_r)]}{(b_z c_z)^3 (c_z - b_z)} &= 0 \\ 2 \frac{d^2 g_z}{dr^2} (-1 + a_z) a_z^3 (b_z c_z)^4 - g_r (-1 + a_r) a_r^3 (b_r c_r)^4 + \frac{g_z}{r^2} (3(-1 + a_z) a_z^3 b_z^2 c_z^2 (b_z - c_z)^2) &= 0 \\ 2 \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{dg_\Phi^I}{dr} \right) (-1 + a_\Phi) a_\Phi^3 (b_\Phi c_\Phi)^4 \\ - \frac{6}{r^2} |z|^2 g_\Phi^I (-1 + a_{\Phi,z}) a_\Phi^3 (c_\Phi^4 b_\Phi^2 + b_\Phi^4 c_\Phi^2) &= 0. \end{aligned} \quad (20)$$

By eliminating $\hat{\mathcal{A}}_r^0$ from the first two equations, one gets the following differential equation for g_z :

$$\begin{aligned} \frac{d^2 g_z}{dr^2} + \frac{dg_r}{dr} \frac{(-1 + a_r) a_r^3 (b_r c_r)^4 \sqrt{3} (-1 + a_z) a_z^3 [(b_z c_z)^3 (c_z - b_z)]}{4(-1 + a_z)^2 a_z^6 (b_z c_z)^4 [b_r^3 c_r^3 \sqrt{3} (2c_r - b_r)]} \\ + \frac{g_z}{r^2} (3(-1 + a_z) a_z^3 \frac{(b_z - c_z)^2}{2b_z^2 c_z^2}) &= 0 \end{aligned} \quad (21)$$

The equation (21) is of the form

$$r^2 y''(r) + \alpha r^2 y'(r) + \beta y(r) = 0, \quad (22)$$

the solution to which (using maple) is given by:

$$\begin{aligned} y(r) &= C_1 \sqrt{r} I_{\frac{1}{2}\sqrt{1-4\beta}} \left(\frac{1}{2} \alpha r \right) e^{-\frac{1}{2} \alpha r} \\ &+ C_2 \sqrt{r} I_{-\frac{1}{2}\sqrt{1-4\beta}} \left(\frac{1}{2} \alpha r \right) e^{-\frac{1}{2} \alpha r} \end{aligned} \quad (23)$$

for $\frac{1}{2}\sqrt{1-4\beta} \notin \mathbf{Z}$ and

$$\begin{aligned} y(r) &= C'_1 \sqrt{r} I_{\frac{1}{2}\sqrt{1-4\beta}} \left(\frac{1}{2} \alpha r \right) e^{-\frac{1}{2} \alpha r} \\ &+ C'_2 \sqrt{r} K_{\frac{1}{2}\sqrt{1-4\beta}} \left(\frac{1}{2} \alpha r \right) e^{-\frac{1}{2} \alpha r} \end{aligned} \quad (24)$$

for $\frac{1}{2}\sqrt{1-4\beta} \in \mathbf{Z}$. Lets consider the two cases separately.

For the former, to make sure that the solution goes to zero as $r \rightarrow \infty$, one sets:

$$C_1 = -C_2. \quad (25)$$

Even though this is a singular solution (at $r = 0$), as we shall see, it is a finite energy solution.

The derivative of the solution involving $I_{\pm\nu}(\alpha r)$ is given by:

$$\begin{aligned} & \frac{d}{dr} \left(e^{-\frac{\alpha r}{2}} \sqrt{r} \left(I_{\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - I_{-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right) \right) \\ &= \frac{e^{-\frac{\alpha r}{2}}}{4\sqrt{r}} \left(\alpha r I_{-1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) + \alpha r I_{1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) + 2(-1 + \alpha r) I_{-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right. \\ & \quad \left. - \alpha r I_{-1-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - \alpha r I_{1-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - 2(-1 + \alpha r) I_{\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right). \quad (26) \end{aligned}$$

For the purpose of getting finite energy, as we will very shortly see, one requires to impose that $r^2 \left(\frac{dy(r)}{dr} \right)^2 \rightarrow 0$ as $r \rightarrow \infty$, and that it is finite at $r = 0$. Using the series expansions around $r = 0$ and the asymptotic expansions for $I_\nu(r)$, the former is satisfied identically, and the latter is satisfied if one imposes:

$$\sqrt{1-4\beta} < 1 \Leftrightarrow 0 < \beta < \frac{1}{4}. \quad (27)$$

The solution for $g_\Phi^I(r)$ that one gets is:

$$g_\Phi^I(r) = 0. \quad (28)$$

Hence, the final result for the $U(1)$ components of the Higgs and gauge fields, in the neighborhood of the South Pole (approaching it along ϕ close to $\frac{\pi}{2}$) are:

$$\begin{aligned} \hat{A}_z^0(r, z) &= i\epsilon f_z(\Theta) \sqrt{r} e^{-\frac{1}{2}\alpha r} \left(I_{\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{1}{2}\alpha r\right) - I_{-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{1}{2}\alpha r\right) \right) \\ &\times \text{Det}[(P_0 - a_z \mathbf{1}_4)(P_1 - M_r)(P_2 - M_r)] + \text{cc}; \\ \hat{A}_r^0 &= i\epsilon f_r(\Theta) \\ &\times \frac{e^{-\frac{\alpha r}{2}}}{4\sqrt{r}} \left(\alpha r I_{-1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) + \alpha r I_{1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) + 2(-1 + \alpha r) I_{-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right. \\ &\quad \left. - \alpha r I_{-1-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - \alpha r I_{1-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - 2(-1 + \alpha r) I_{\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right) \\ &\times (z \text{Det}[(P_0 - a_z \mathbf{1}_4)(P_1 - M_r)(P_2 - M_r)] + \text{cc}); \\ \hat{\Phi}^0 &= 0, \quad (29) \end{aligned}$$

where

$$\begin{aligned}
\beta &\equiv (3(-1 + a_z)a_z^3 \frac{(b_z - c_z)^2}{2b_z^2 c_z^2}) < 0; \\
\alpha &\equiv \frac{(-1 + a_r)a_r^3(b_r c_r)^4 \sqrt{3}(-1 + a_z)a_z^3[(b_z e_z)^3(c_z - b_z)]}{4(-1 + a_z)^2 a_z^6(b_z c_z)^4[b_r^3 c_r^3 \sqrt{3}(2c_r - b_r)]}; \\
0 &< a_z < 1; \quad c_{z,r} = e_{z,r}; \quad b_{z,r} = f_{z,r}.
\end{aligned} \tag{30}$$

Lets consider the energy of the above solution. It is given by (we follow the conventions of [1]):

$$\begin{aligned}
E &= -\frac{1}{4\pi} \int r^2 dr \frac{dz d\bar{z}}{(1 + |z|^2)^2} \text{Tr} \left[(\hat{D}_i * \hat{\Phi})^2 + \frac{1}{2} (\hat{F}_{ij})^2 * \right] \\
&= E_{SU(4)} + E_{U(1)}.
\end{aligned} \tag{31}$$

But, in the neighborhood of the South Pole, where one can drop the $*$ product,

$$\begin{aligned}
E_{U(1)} &= - \int r^2 dr \frac{dz d\bar{z}}{(1 + |z|^2)^2} \left[(\partial_r \hat{\mathcal{A}}^0_z - \partial \hat{\mathcal{A}}^0_r)^2 + (\partial_r \hat{\mathcal{A}}^0_z + \bar{\partial} \hat{\mathcal{A}}^0_r)^2 \right. \\
&\quad \left. + ((\partial + \bar{\partial}) \hat{\mathcal{A}}^0_z)^2 \right] \geq 0.
\end{aligned} \tag{32}$$

Hence, in the neighborhood of the South Pole, even though one is dealing with non-BPS solutions, one gets the following inequality:

$$E_{U(4)} \geq E_{SU(4)}, \tag{33}$$

with saturation of the inequality occuring for null $U(1)$ component. Substituting (29) in the $Im \ z \rightarrow \infty$ limit into (32), and integrating in the neighborhood of the South Pole as described earlier, i.e., around $\phi = \frac{\pi}{2}$ and $\theta \in [\pi - \epsilon, \pi]$, $\lim_{\epsilon \rightarrow 0}$, one gets an infinitesimal result, provided one gets a finite result from the r integration. The r -integral should converge to get a finite-energy solution. One sees that one has to evaluate:

$$\begin{aligned}
E_{U(1)} \sim \int_0^\infty dr r^2 \frac{e^{-\alpha r}}{16r} &\left(\alpha r I_{-1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) + \alpha r I_{1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) + 2(-1 + \alpha r) I_{-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right. \\
&\left. - \alpha r I_{-1-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - \alpha r I_{1-\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - 2(-1 + \alpha r) I_{\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right)^2.
\end{aligned} \tag{34}$$

We have not been able to evaluate the above integral. But the integrand for $0 < \beta < \frac{1}{4}$ is analytic for all values of $r \in [0, \infty)$, and it vanishes at $r = 0$ and $r \rightarrow \infty$.

We hence assume that it is convergent corresponding to a finite energy solution. We have verified the same numerically using mathematica for several values of $\beta < \frac{1}{4}$.

Let us now consider the case $\frac{1}{2}\sqrt{1-4\beta} \in \mathbf{Z}$. In that case in order to get a solution that vanishes as $r \rightarrow \infty$, one sets $C'_1 = 0$ in (24).

One then requires:

$$E_{U(1)} \sim \int_0^\infty dr r^2 \frac{e^{-\alpha r}}{16r} \left(-\alpha r K_{-1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - 2(-1+\alpha r) K_{\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) - \alpha r K_{1+\frac{1}{2}\sqrt{1-4\beta}}\left(\frac{\alpha r}{2}\right) \right)^2 < \infty. \quad (35)$$

By using the series expansion of $K_\nu(x)$ about $x = 0$ (See [8]), one sees that one can not find a $\beta : \frac{1}{2}\sqrt{1-4\beta} \in \mathbf{Z}$ that will satisfy (35) because of the singular nature of the integrand at $r = 0$.

Now, lets turn to calculating the commutative Higgs and gauge field: $\Phi^I, A_{r,z,\bar{z}}$. As we have thus far continued working to all orders in the noncommutativity parameter Θ , one hence requires an all-order (in Θ) Seiberg-Witten map [9, 10]. There is a conjecture for that as given in [10]. We will not explicitly construct the SW map, but conjecture that in the neighborhood of the negative x^3 axis (or the neighborhood of the South Pole of the Riemann Sphere, approaching it the way described in the paper), the noncommutative hatted fields are the same as the commutative unhatted fields. Noting that as $Imz \rightarrow \infty$, $\hat{A}_{z,\bar{z}}^{SU(4)}, \partial, \bar{\partial}\hat{\Phi}^{SU(4)}, \partial\hat{A}_z^{U(1)}, \partial^2\hat{A}_r^{U(1)} \rightarrow 0$, and dropping terms of $O(\epsilon^2)$ in $\hat{A}_{r,z}^{U(1)}$ in equations (3.3), (3.4) and (3.7) for the Higgs and equations (A3), (A6) and (A8) for the gauge field of [12], the conjecture can be explicitly verified to $O(\Theta^2)$. Hence, in the neighborhood of the South Pole, approaching it along an azimuthal angle close to $\frac{\pi}{2}$

$$\begin{aligned} \hat{\Phi} &= \hat{\Phi}^a T^a = \Phi^a T^a; \\ \hat{A}_i &= \hat{A}_i^a T^a + \hat{A}_i^0 \mathbf{1}_4 = A_i^a T^a + A_i^0 \mathbf{1}_4. \end{aligned} \quad (36)$$

The electric charge is then the same as that given in [1]. The magnetic charge will receive a $U(1)$ contribution proportional to $(\partial + \bar{\partial})A_z(z, \bar{z}, r \rightarrow \infty)$, which using (29), vanishes. Hence, the noncommutative dyon in the neighborhood of the South Pole, carries the same electric and magnetic charges as the commutative dyon of [1]. Their energy will be more than that of the commutative dyons in the neighborhood of the South Pole, though by an infinitesimal amount. As the equality of the commutative and noncommutative electric and magnetic charges is valid only in the neighborhood of the South Pole, these noncommutative dyonic solutions are likely to be stable.

4 Summary and Discussion

To summarize, we have shown that noncommutative (with $r - z$ and $r - \bar{z}$ noncommutativity) $U(4)$ SYM possesses static dyonic solutions in the neighborhood of the South Pole of the Riemann sphere or equivalently the negative x^3 axis, that either are the same as the static commutative $SU(4)$ dyonic solutions of [1] (corresponding to *decoupling of a null $U(1)$ from the $SU(4)$*) or, have the same electric and magnetic charges as the commutative dyons and differ infinitesimally in their energy (corresponding to *decoupling of a non-null $U(1)$ from the $SU(4)$*), to *all orders in the noncommutativity parameter*. We have thus shown the decoupling of the $U(1)$ sector, both for null and infinitesimal $U(1)$ components in the neighborhood of the negative x^3 axis. This decoupling to $O(\Theta)$ for the whole of space, was also seen in the context of noncommutative monopole solutions in SYM in [3]. The noncommutative hatted fields are the same as the commutative unhatted fields (we conjecture to all orders in the noncommutativity parameter and show explicitly up to $O(\Theta^2)$). Thus, we see that the neighborhood of the negative x^3 axis, acts as “denoncommutatifier”. It will be interesting to study the same problem using noncommutative ADHMN construction (See [14] and references therein), and to be able to establish the existence of such solutions using techniques of [15] using Morse theory. Also, finding similar solutions with $z - \bar{z}$ noncommutativity will be nice.

Acknowledgements

We thank A.Kumar for useful discussions. We also thank K.L.Panigrahi, R.R.Nayak, S.S.Pal and G.Bhattacharya for bringing to our notice references [1], [10] and [5].

Appendix A

In this appendix, we summarize the asymptotic (w.r.t to z) expressions for $A_{r,z}^{U(1)}$ and some expressions involving them necessary to obtain the $U(1)$ components of the gauge field and Higgs. We have used mathematica for arriving at the expressions given in this appendix.

Using (10), one can explicitly write out the expressions for $\hat{\phi}^0, \hat{A}_z^0 = \text{Det}[(P_0 - a_{\Phi,z}\mathbf{1}_4)(P_1 - M_{\Phi,z}).(P_2 - M_{\Phi,z})] + \text{cc}$ and $\hat{A}_r^0 = z\text{Det}[(P_0 - a_r\mathbf{1}_4)(P_1 - M_r).(P_2 - M_r)] + \text{cc}$, using mathematica, but because of the length of the expressions, we will give only their asymptotic forms.

One gets as $\text{Im}z \rightarrow \infty$:

$$\hat{\phi}^0, \hat{A}_z^0 \sim 2(-1 + a_{\Phi,z})a_{\Phi,z}^3(b_{\Phi,z}c_{\Phi,z}e_{\Phi,z}f_{\Phi,z})^2. \quad (\text{A1})$$

Similarly, as $z \rightarrow \infty$, one gets:

$$\hat{A}_r^0 \sim 2(-1 + a_r)a_r^3(b_rc_re_rf_r)^2 Re z. \quad (A2)$$

Hence, we see that if $z \rightarrow \infty$ as $z = (Rez)_0 + i\infty$, where $(Rez)_0$ is finite, then $\hat{A}_r$ is non-singular, else it is singular.

One sees that as $z \rightarrow \infty$,

$$\partial \hat{A}_r^0 \sim (-1 + a_r)a_r^3(b_re_rc_rf_r)^2. \quad (A3)$$

For the purpose of calculation of energy and imposing it to be finite, one requires to calculate $\bar{\partial} \hat{A}_r^0$, and its asymptotic expression is given by:

$$\bar{\partial} \hat{A}_r^0 \sim (-1 + a_r)a_r^3 b_r^2 c_r^2 e_r^2 f_r^2 \quad (A4)$$

One also needs to evaluate $\partial \bar{\partial} \hat{A}_r^0$ as $z \rightarrow \infty$, and is given by:

$$\partial \bar{\partial} \hat{A}_r^0 \sim (-1 + a_r)a_r^3 \frac{b_r^2 c_r e_r^2 f_r (\sqrt{3}c_r - \sqrt{3}f_r)\bar{z}^2 + \sqrt{3}(b_re_rc_r)^2 f_r z^2}{|z|^4} \quad (A5)$$

Also, one one sees that:

$$\partial \bar{\partial}(\hat{\phi}^0, \hat{A}_z^0) \sim -6(-1 + a_{\Phi,z})a_{\Phi,z}^3 \frac{(((c_{\Phi,z}e_{\Phi,z})^2 b_{\Phi,z} f_{\Phi,z} + (b_{\Phi,z} f_{\Phi,z})^2 c_{\Phi,z} e_{\Phi,z}))}{|z|^2}. \quad (A6)$$

For the purpose of solving for the $U(1)$ components of the gauge field, and for getting the magnetic charge of the dyon, one needs to evaluate $(\partial + \bar{\partial})\hat{A}_z^0$, and as $z \rightarrow \infty$, one gets:

$$(\partial + \bar{\partial})\hat{A}_z^0 \sim \frac{b_z c_z^2 e_z f_z^2 (\sqrt{3}b_z - \sqrt{3}e_z)\bar{z}^2 + (c_z f_z b_z^2 (-e_z f_z \sqrt{3}e_z + c_z e_z \sqrt{3}e_z) - 3c_z f_z b_z e_z c_z e_z)z^2}{|z|^4} \quad (A7)$$

Two other quantities that one needs to calculate are: $(\partial - \bar{\partial})(\hat{\Phi}, \hat{A}_z)$ and $\partial \left((1 + |z|^2)(\partial + \bar{\partial})(\hat{\Phi}, \hat{A}_z) \right)$. The former as $z \rightarrow \infty$, is given by:

$$(\partial - \bar{\partial})(\hat{\phi}^0, \hat{A}_z^0) \sim \frac{\sqrt{3}(-1 + a_i)a_i^3 \left[(b_i e_i)^2 c_i f_i (c_i - f_i)\bar{z} + (-b_i c_i^2 e_i^2 f_i (f_i - b_i) + e_i b_i^2 c_i f_i^2 (c_i - e_i))z \right] (z - \bar{z})}{|z|^4}, \quad (A8)$$

The latter as $z \rightarrow \infty$ is given by:

$$\begin{aligned} & \partial \left((1 + |z|^2)^2 (\partial + \bar{\partial})(\phi^0, \hat{\mathcal{A}}_z^0) \right) \sim \\ & -2(-1 + a_i) a_i^3 \left(\sqrt{3} [c_i^2 e_i^2 b_i f_i (-f_i + b_i) + b_i^2 f_i^2 c_i e_i (c_i - e_i)] z - (3b_i f_i (c_i e_i)^2 + 3c_i e_i (b_i f_i)^2) \right. \\ & \left. + \frac{3b_i c_i e_i f_i (b_i e_i + c_i f_i) \bar{z}^4}{|z|^4} \right) \end{aligned} \quad (\text{A9})$$

References

- [1] T. Ioannidou and P. M. Sutcliffe, *Non-BPS string junctions and dyons in $N = 4$ super-Yang-Mills*, Phys. Lett. B **467**, 54 (1999) [hep-th/9907157]; *Non-Bogomolny $SU(N)$ BPS monopoles*, Phys. Rev. D **60**, 105009 (1999) [hep-th/9905169].
- [2] T. Ioannidou and P. M. Sutcliffe, *Monopoles and harmonic maps*, J. Math. Phys. **40**, 5440 (1999) [hep-th/9903183].
- [3] K. Hashimoto, H. Hata and S. Moriyama, *Brane configuration from monopole solution in non-commutative super Yang-Mills theory*, JHEP **9912**, 021 (1999) [hep-th/9910196].
- [4] H. Hata and S. Moriyama, *String junction from non-commutative super Yang-Mills theory*, JHEP **0002**, 011 (2000) [hep-th/0001135].
- [5] K. Matsubara, *Restrictions on gauge groups in noncommutative gauge theory*, Phys. Lett. B **482**, 417 (2000) [hep-th/0003294].
- [6] B. Jurco, L. Moller, S. Schraml, P. Schupp and J. Wess, *Construction of non-Abelian gauge theories on noncommutative spaces*, hep-th/0104153.
- [7] T. Ioannidou, B. Piette and W. J. Zakrzewski, *Low energy states in the $SU(N)$ Skyrme models*, hep-th/9811071.
- [8] *Bateman Manuscript Project, Higher Transcendental Functions, Vol II*, A. Erdelyi (editor), McGraw-Hill Book Company (NY), 1953.
- [9] N. Seiberg and E. Witten, *String theory and noncommutative geometry*, JHEP **9909**, 032 (1999) [hep-th/9908142].
- [10] H. Liu, **-Trek II: *n operations, open Wilson lines and the Seiberg-Witten map*, hep-th/0011125.

- [11] T. Asakawa and I. Kishimoto, *Comments on gauge equivalence in noncommutative geometry*, JHEP **9911**, 024 (1999) [hep-th/9909139].
- [12] S. Goto and H. Hata, *Noncommutative monopole at the second order in theta*, Phys. Rev. D **62**, 085022 (2000) [hep-th/0005101].
- [13] E. S. Na, *Superpartner Solutions of a BPS Monopole in Noncommutative Space*, hep-th/0106278.
- [14] D. Bak, *Deformed Nahm equation and a noncommutative BPS monopole*, Phys. Lett. B **471**, 149 (1999) [hep-th/9910135].
- [15] C. H. Taubes, *Monopoles And Maps From $S(2)$ To $S(2)$: The Topology Of The Configuration Space*, Commun. Math. Phys. **95** (1984) 345; *The Existence Of A Nonminimal Solution To The $SU(2)$ Yang-Mills Higgs Equations On R^3* , Commun. Math. Phys. **86**, 257 (1982).

